
MONOCLONAL ANTIBODY TO XYLOGLUCAN (LM25) (RAT IgM, KAPPA LIGHT)

FOR RESEARCH PURPOSES ONLY

Cat. No. 600-8

SPECIFICITY

This antibody recognizes xyloglucan. Generated using a neoglycoprotein incorporating the xylosylated/galactosylated oligosaccharides from tamarind xyloglucan (XXLG and XLLG). It recognizes a range of xyloglucan-specific oligosaccharide motifs in several species.

It recognizes xyloglucan from Tamarind seed (Megazyme P-XYGLN).

This antibody can be used with second stage anti-rat antibodies labelled with gold, FITC, HRP etc.

USE

Reconstitute in 150 μ L of 120mM NaCl.

When reconstituted, the antibody concentration is 1mg/mL in 10mM phosphate buffer (pH 7.2), 0.8% sodium chloride.

In dot blot assays we found the 1mg/mL solution to be 10 to 100x more sensitive than the original *Plant Probes* generated supernatant.

STORAGE

This freeze-dried product has been shipped at ambient temperature; however, it should be stored for extended periods at less than 4°C, with desiccant.

Once reconstituted, store at 4°C short term, or at -20°C in aliquots longer term. Do not re-freeze after thawing. Once thawed, store at 4°C.

REFERENCES

Sims, I.M., Munro, S.L.A., Currie, G., Criak, D., Bacic, A. (1996). *Carbohydrate Research*, 293: 147-172

Pedersen, H.L., Fangel, J.U., McCleary, B., Ruzanski, C., Rydahl, M.G., Ralet, M.C., Farkas, V., von Schantz, L., Marcus, S.E., Andersen, M.C. and Field, R. (2012). *Journal of Biological Chemistry*, 287(47): 39429-39438.

Ruprecht, C., Bartetzko, M.P., Senf, D., Dallabernadina, P., Boos, I., Andersen, M.C., Kotake, T., Knox, J.P., Hahn, M.G., Clausen, M.H. and Pfrengle, F. (2017). *Plant Physiology*, 175: 1094-1104.

(Under licence from University of Leeds, UK)

Biosupplies Australia Pty Ltd

A.B.N. 62 006 209 034

P.O. Box 187, La Trobe University, Bundoora, Victoria 3083, AUSTRALIA
www.biosupplies.com.au