

### **Citations for Cat#100-1: Aniline blue fluorochrome**

Stone, BA, Evans, NA, Bonig, I, Clarke, AE (1984). The application of sirofluor, a chemically defined fluorochrome from aniline blue for the histochemical detection of callose. *Protoplasma* 122 (3), 191-195.

### **Metal toxicity**

Wissemeier, AH, Klotz, F, Horst, WJ (1987). Aluminum induced callose synthesis in roots of soybean (glycine-max L). *Journal of Plant Physiology* 129 (5), 487-492.

### **Plant cell biology**

Ohana, P, Benziman, M, Delmer, DP (1993). Stimulation of callose synthesis in vivo correlates with changes in intracellular-distribution of the callose synthase activator beta-furfuryl-beta-glucoside. *Plant Physiology* 101 (1), 187-191.

Webb, MC, Williams, EG (1988). Effects of temperature, light, nutrients and carbon-dioxide on the strength of the self-incompatibility response in detached flowers of *Lycopersicon-Peruvianum*. *Annals Of Botany* 61 (4), 395-404.

Hayashi, T, Yoshida, K, (1988). Cell expansion and single-cell separation induced by colchicine in suspension-cultured soybean cells. *Proceedings Of The National Academy Of Sciences Of The United States Of America* 85 (8), 2618-2622.

Foissner, I (1988). Chlortetracycline-induced formation of wall appositions (callose plugs) in internodal cells of *Nitella-Flexilis* (Characeae). *Journal Of Phycology* 24 (4), 458-467.

Suzuki, K, Amino, S, Takeuchi, Y, et al. (1990). Differences in the composition of the cell-walls of 2 morphologically different lines of suspension-cultured *Catharanthus-Roseus* cells. *Plant And Cell Physiology* 31 (1), 7-14.

Bayles CJ, Ghemawat MS, Aist JR (1990). Inhibition by 2-deoxy-d-glucose of callose formation, papilla deposition, and resistance to powdery mildew in an ml-o barley mutant. *Physiological And Molecular Plant Pathology* 36 (1), 63-72.

Webb, MC, Williams, EG (1988). Effects of temperature, light, nutrients and carbon-dioxide on the strength of the self-incompatibility response in detached flowers of *Lycopersicon-Peruvianum*. *Annals Of Botany* 61 (4), 395-404.

Bernatzky, R, Anderson, MA, Clarke, AE (1988). Molecular-genetics of self-incompatibility in flowering plants. *Developmental Genetics* 9 (1), 1-12.

Gaude, T, Dumas, C (1987). Molecular and cellular events of self-incompatibility. *International Review Of Cytology-A Survey Of Cell Biology* 107, 333-366.

Yim, KO, Bradford, KJ (1998). Callose deposition is responsible for apoplastic semipermeability of the endosperm envelope of muskmelon seeds. *Plant Physiology* 118 (1), 83-90.

Nickle, TC, Meinke, DW (1998). A cytokinesis-defective mutant of *Arabidopsis* (cyt1) characterized by embryonic lethality, incomplete cell walls, and excessive callose accumulation. *Plant Journal* 15 (3), 321-332.

McConchie, CA, Knox, RB (1989). Pollen-stigma interaction in the seagrass *Posidonia australis*. *Annals of Botany* 63 (2): 235-248

Jacobs, AK, Lipka, V, Burton, RA, et al. (2003). An *Arabidopsis* callose synthase, GSL5, is required for wound and papillary callose formation. *Plant Cell* 15 (11), 2503-2513.

Freialdenhoven, A, Peterhansel, C, Kurth, J, et al. (1996). Identification of genes required for the function of non-race-specific mlo resistance to powdery mildew in barley. *Plant Cell* 8 (1), 5-14.

Gorshkova, EN, Erokhina, TN, Stroganova, TA, et al. (2003). Immunodetection and fluorescent microscopy of transgenically expressed hordeivirus TGBp3 movement protein reveals its association with endoplasmic reticulum elements in close proximity to plasmodesmata. *Journal of General Virology* 84: 985-994.

### **Plant histochemistry**

Serpe, MD, Muir, AJ, Keidel, AM (2001). Localization of cell wall polysaccharides in nonarticulated laticifers of *Asclepias speciosa* Torr. *Protoplasma* 216 (3-4), 215-226.

Gane, AM, Weinhandl, JA, Bacic, A, et al. (1994). Histochemistry and composition of the cell-walls of styles of *Nicotiana-alata* link et otto. *Planta* 195 (2), 217-225.

Rougier, M, Jnoud, N, Said, C, et al. (1991). Male gametophyte development and formation of the male germ unit in populus-deltoides following compatible pollination. *Protoplasma* 162 (2-3), 140-150.

Rae, AL, Harris, PJ, Bacic, A, et al. (1985). Composition of the cell-walls of *Nicotiana alata* link et otto pollen tubes. *Planta* 166 (1), 128-133.

Meyberg, M (1988). Cyto-chemistry and ultrastructure of the mucilage secreting trichomes of nymphaeoides-peltata (menyanthaceae). *Annals of Botany* 62 (5), 537-547.

White J (1990). Pollen development in actinidia-deliciosa var deliciosa - histochemistry of the microspore mother cell-walls. *Annals of botany* 65 (3), 231-239.

Anderson, MA, Harris, PJ, Bonig, I, et al. (1987). Immunogold localization of alpha-l-arabinofuranosyl residues in pollen tubes of *Nicotiana glauca* Link et Otto. *Planta* 171 (4), 438-442.

Alche, JD, Rodriguez-Garcia, MI (1997). Fluorochromes for detection of callose in meiocytes of olive (*Olea europaea* L.) *Biotechnic and Histochemistry* 72 (6), 285-290.

Pline, WA, Edmisten, KL, Oliver, T, et al. (2002). Use of digital image analysis, viability stains, and germination assays to estimate conventional and glyphosate-resistant cotton pollen viability. *Crop Science* 42 (6), 2193-2200.

Yamamoto, M, Nozaki, H, Miyazawa, Y, et al. (2003). Relationship between presence of a mother cell wall and speciation in the unicellular microalga *Nannochloris* (Chlorophyta). *Journal of Phycology* 39 (1), 172-184.

### ***In vitro* synthesis products**

Kudlicka, K, Brown, RM (1997). Cellulose and callose biosynthesis in higher plants .1. Solubilization and separation of (1-3)- and (1-4)-beta-glucan synthase activities from mung bean. *Plant Physiology* 115 (2), 643-656.

Hayashi, T, Read, SM, Bussell, J, et al. (1987). UDP-glucose-(1-3)-beta-glucan synthases from mung bean and cotton - differential-effects of Ca<sup>2+</sup> and Mg<sup>2+</sup> on enzyme properties and on macromolecular structure of the glucan product. *Plant Physiology* 83 (4), 1054-1062.

Ahlborn, B, Werner, D (1992). 1,3-beta-glucan synthase on the peribacteroid membrane (symbiosome membrane) from soybean root-nodules. *Physiological and molecular plant pathology* 40 (5), 299-314.

### **Fungal exo-polysaccharides**

Michel, M, Seviour, RJ, Pethica, LM (1987). Exocellular polysaccharide production by isolates of *Epicoccum purpurascens*. *Biotechnology letters* 9 (10), 741-744.

Stasinopoulos, SJ, Seviour, RJ (1989). Exopolysaccharide formation by isolates of *Cephalosporium* and *Acremonium*. *Mycological research* 92, 55-60.

### **Plant-Microorganism Interactions**

Aist JR, Gold RE, Bayles CJ, et al. (1988). Evidence that molecular-components of papillae may be involved in ml-o resistance to barley powdery mildew. *Physiological and molecular plant pathology* 33 (1), 17-32.

Drews, G, Ziser, K, Schrockvictor, U, et al. (1988). Cellular-responses of soybean to virulent and avirulent strains of *Pseudomonas syringae* pv *glycinea*. *European journal of cell biology* 46 (3), 369-375.

Bonhoff, A, Rieth, B, Golecki, J, et al. (1987). Race cultivar-specific differences in callose deposition in soybean roots following infection with *Phytophthora megasperma* f-sp *glycinea*. *Planta* 172 (1), 101-105.

Jordan, CM, Endo, RM, Jordan, LS (1988). Penetration and colonization of resistant and susceptible apium-graveolens by *Fusarium oxysporum* f-sp *apii* race-2 - callose as a structural response. *Canadian journal of botany-revue canadienne de botanique* 66 (12), 2385-2391.

Bretschneider, KE, Gonella, MP, Robeson, DJ (1989). A comparative light and electron microscopical study of compatible and incompatible interactions between *Xanthomonas campestris* pv *campestris* and cabbage (*Brassica oleracea*). *Physiological and molecular plant pathology* 34 (4), 285-297.

Mcgovern, RJ, Horst, RK, Israel, HW (1989). *Chrysanthemum* phloem necrosis - symptomatology and histopathology. *Canadian journal of botany-revue canadienne de botanique* 67 (4), 1014-1023.

Mcgovern, RJ, Horst, RK, Israel, HW (1989). *Chrysanthemum* phloem necrosis - detection by epifluorescence microscopy. *Canadian journal of botany-revue canadienne de botanique* 67 (4), 1024-1031.

Lotan, T, Ori, N, Fluhr, R (1989). Pathogenesis-related proteins are developmentally regulated in tobacco flowers. *Plant cell* 1 (9), 881-887.

Cohen, Y, Eyal, H, Hanania, J (1990). Ultrastructure, autofluorescence, callose deposition and lignification in susceptible and resistant muskmelon leaves infected with the powdery mildew fungus *Sphaerotheca fuliginea*. *Physiological and molecular plant pathology* 36 (3), 191-204.

Golotte, A, Gianinazziperson, V, Giovannetti, M, et al. (1993). Cellular-localization and cytochemical probing of resistance reactions to arbuscular mycorrhizal fungi in a locus a myc-mutant of *Pisum sativum* l. *Planta* 191 (1), 112-122.

Nicholas, RO, Williams, DW, Hunter, PA (1994). Investigation of the value of beta-glucan-specific fluorochromes for predicting the beta-glucan content of the cell-walls of zoopathogenic fungi. *Mycological research* 98, 694-698.

## **Yeast morphogenesis**

Kippert F, Lloyd D (1995). The aniline blue fluorochrome specifically stains the septum of both live and fixed schizosaccharomyces-pombe cells. *FEMS microbiology letters* 132 (3), 215-219.

## **Other**

Hough, T, Bernhardt, P, Knox, RB, Williams, EG (1985). Applications of fluorochromes to pollen biology. II. The DNA probes ethidium bromide and Hoechst 33258 in conjunction with the callose-specific aniline blue fluorochrome. *Stain Technol.* 60(3),155-62.

Yim. K-O, Bradford, KJ (1998). Callose Deposition Is Responsible for Apoplastic Semipermeability of the Endosperm Envelope of Muskmelon Seeds. *Plant Physiology* 118, 83-90.

Cholewa, E, Griffith, M (2004). The unusual vascular structure of the corm of *Eriophorum vaginatum*: implications for efficient retranslocation of nutrients. *Journal of Experimental Botany* 55 (397), 731-741.